



Pressure Injuries

Information for Patients, Residents and Carers

What is a Pressure Injury?

A pressure injury is a localised injury to the skin and/or underlying tissue due to unrelieved **pressure, friction, shear,** or **moisture** or a combination of these factors. Pressure ulcer, decubitus ulcer and bedsore are other terms used for this problem.

While pressure injuries can appear insignificant on the surface, under the skin damage can occur which can spread into surrounding muscles and bones, leading to pain, possible infection, scarring and a delayed recovery.

Risk is increased with:

- reduced mobility
- altered mental state
- poor nutrition
- incontinence
- reduced sensation
- ill health

What to look for?

- Red/Purple/Blue skin
- Blisters
- Swelling
- Patches of dry skin and hard skin
- Shiny areas

- Warm or cool areas
- Cracks, calluses, wrinkles
- Localised pain

Where to look?

Pressure injuries can occur anywhere there is continued pressure to the skin but usually over bony areas – especially the heels, elbows, hips, buttocks and tail bone. Pressure injuries can also be caused by medical devices such as splints, compression hosiery and masks.



What you can do

- ✓ Change position frequently to prevent pressure damage
- ✓ Use soap-free bathing and cleanser products to help protect your skin from drying out
- ✓ Moisturise your skin twice daily
- ✓ Protect skin exposed to friction, pat skin dry and avoid rubbing
- ✓ Use pillows and foam wedges to protect bony areas
- ✓ Eat a balanced diet including two servings of fruit and five servings of vegetables each day
- ✓ Drink 6 to 8 glasses of water every day
- ✓ Check your skin regularly and contact your health professional or the Wound Innovations team if you have any soreness, reddened, blistered or broken skin.

Remember – Keep on moving! Look after your skin and eat a healthy diet!

What you should NOT do!

- ✗ Do not massage or rub the skin as this causes friction
- ✗ Avoid sitting in a chair or wheelchair for long periods of time
- ✗ Do not use foam rings or donuts as these have been found to cause more pressure
- ✗ Avoid products that irritate the skin. Try unperfumed lotions and avoid talcum powder.

Contact Information

Wound Innovations: [1300 224 034](tel:1300224034)

reception@woundinnovations.com.au www.woundinnovations.com.au

References

Wound Innovations. (2018). Assessment and Management of Pressure Injuries Online Education Module. QLD, Australia: Wound Innovations.

European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel and Pan Pacific Pressure Injury Alliance. (2019). Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline. The International Guideline. Emily Haesler (Ed.). EPUAP/NPIAP/PPPIA.

National Health and Medical Research Council. (2013). Australian Dietary Guidelines [National Guidelines]. Canberra, Australia: National Health and Medical Research Council

N.I.C.E. (2014). Pressure Ulcers: prevention and management [Clinical Guideline]. Retrieved from: www.nice.org.uk

DISCLAIMER: This document is intended to be used as a guide of general nature, having regard to general circumstances. It cannot be comprehensive and is intended as a guide only. This document reflects information available at the time of its preparation, but its currency should be determined having regard to other available information. The information presented should not be relied on as a substitute for medical advice, independent judgement or proper assessment by a doctor, with consideration of the particular circumstances of each case and individual needs. Wound Innovations disclaims all liability to users of the information provided.